

SF (MARGINS- related)	Collaborative Research: Hf-Nd Isotopic and Trace-Element Geochemistry of Globally Subducting Sediments	
	Katherine Kelley & Terry Plank, Boston Univ.; John Ludden, CNRS; Hubert Staudigel, Scripps Inst. of Oceanography; Vervoort	
	2/28/2005 – 2/28/2005	OCE 01-37110

Many studies have argued, based partly on Pb isotopic evidence, that recycled subducted slabs reside in the mantle source of ocean island basalts. Such models, however, have remained largely untested against actual subduction zone inputs, due to the scarcity of comprehensive measurements of both radioactive parents (Th and U) and radiogenic daughter (Pb) in altered oceanic crust (AOC). Here, we report comprehensive measurements of U, Th and Pb concentrations in the oldest AOC, ODP Site 801, and consider the effect of subducting this crust on the long-term Pb isotope evolution of the mantle.

- The upper 500 m of AOC at the oldest basement drill site to date (ODP Site 801) shows 5-fold enrichment in U over pristine glass during seafloor alteration, but no net change to Pb or Th.
- Pacific arcs, including the Marianas, define mixing trends toward a single low-Th, high-Pb component derived from the AOC. This AOC-derived component requires 20-fold enrichment in Pb over U during slab dehydration beneath the arc. This output constraint, along with the input constrained from Site 801, enables a mass balance of U, Th and Pb across the Mariana subduction system (see Figure).
- Pb is lost shallow and U deep from subducted AOC, which may be a consequence of the stability of phases that bind these elements during seafloor alteration: U in carbonate and Pb in sulfides.
- Preferred crustal growth rates require U loss at depths beyond the zones of arc and back-arc magmagenesis. Such a U flux could decrease upper mantle k ($^{232}\text{Th}/^{238}\text{U}$) over time, and provide a solution to the "kappa conundrum" (Elliott *et al.*, 1999, EPSL).
- The net effects of alteration and subduction are sufficient to create the Pb isotopic signatures of ocean island basalts, including "himu" end-members (high $^{206}\text{Pb}/^{204}\text{Pb}$).

Figures and Captions




Figure 1: Losses of Pb and U from subducted oceanic crust (excluding sediment) vs. depth in the subduction zone, modeled using inputs and outputs across the Mariana Arc. Grey boxes indicate the full range of possible losses permitted by current estimates of arc crustal growth rates. Heavy

black line traces slab losses at preferred crustal growth rates.

Publications and Presentations

Kelley, K.A., Plank, T., Ludden, J.N. and H. Staudigel (2003) The composition of altered oceanic crust at ODP sites 801 and 1149. *Geochem. Geophys. Geosyst.* 4 (6), doi: 10.1029/2002GC000435.

Kelley, K.A., Plank, T., Ludden, J.N. & Staudigel, H. Subduction Cycling of U, Th, and Pb, soon to be submitted to EPSL.

Farr, L.C., Plank, T., Kelley, K. A. and Alt, J.C. (2001) U Mineral hosts and enrichment processes in altered oceanic crust. *Eos Trans AGU*, 82 (47), Fall Meet. Suppl., Abstract T22C-0926.